# MANAGE HERBIVORES TO PROTECT OR PROMOTE REGENERATION

#### **OBJECTIVE**

Preserve the integrity of the forest.

#### DESCRIPTION

Climate change can increase herbivorous populations that benefit from warmer conditions. Some herbivores can become a stressor in forests and there is potential for them to increase in size and intensity if the above mentioned changes in climate lower winter mortality and allow herbivorous populations to grow. As climate change exacerbates many forest stressors, it will be increasingly important to protect the regeneration of desired species from these animals. At the landscape level, an example of tactic that is sometimes employed to influence deers' habits is to harvest timber in mountain forests to reduce the migration of some species to adjacent coniferous forests, where regeneration is highly vulnerable. Examples of booth-level adaptation tactics include using fences and other barriers, as well as "hiding" more desirable species in a mix of less desirable species.

## **EXPECTED RESULTS**

Preserved forest ecosystem.

#### **RESULT INDICATORS**

Number of plant species preserved

#### **INVOLVED ACTORS**

Environmental agencies and local government.

## **EXPECTED TIMELINE FOR ACTION**

• Short term (1-4 years)

#### **BEST PRACTICES**

- Central Appalachians USA
- Massachusetts USA
- USA
- North America, Europe, Australia and New Zealand

#### CRITICALITIES

Biodiversity conservation; reliable data collection.



## **SCOPE OF THE ACTION**

• Adaptation

#### **TYPE OF PROPOSED ACTIONS**

- Green
- Soft

#### **SECTOR OF ACTION**

- Agriculture / Forests / Land use
- Biodiversity / Conservation of ecosystems
- Other

### **CLIMATE IMPACTS**

- Change or loss of biodiversity
- Other

### **IMPLEMENTATION SCALE**

- Province
- Region / Country

SOURCE

https://www.nrs.fs.fed.us/

